

REPRODUCTION OF THE MAGNOLIA (MAGNOLIACEAE) PLANT IN NAMANGAN CONDITIONS

N. R. Melanova

Annotation:

Magnoliaceae is a family of dicotyledonous plants. Consists of trees and shrubs. The leaves are entire, sometimes lobed, the margins are flat, the flowers are large, straight, bisexual (rarely homogeneous). There are many pollinators. There are 14 genera (about 240 species) in the tropical and subtropical regions of America and Asia. Magnolia and tulip species are cultivated as ornamental plants.

Keywords: magnolia, seeds, cuttings, ornamental, flowers.

Magnolia (Magnolia) is a family of evergreen or deciduous trees belonging to the family Magnolia. East and Southeast Asia. There are about 80 species in Southeast America and Central America. There are 3 species of ornamental plants in Uzbekistan. Hence, the large-flowered Magnolia. Always propagated as a green ornamental tree. Leaves grown in Uzbekistan shed during autumn. It is 10-13 m long. The leaves are elliptic or ovoid, glossy, banded, arranged in series. The flowers are large, white, fragrant, solitary. The leaves contain essential oils, glucosides, alkaloids in the bark and roots. The liquid extract from the leaves is used to lower blood pressure.

The beauty of the leaves, the fragrant flowers, the uniqueness of the fruit, the magnolia is a solitary, as well as ornamental flowering evergreen or deciduous trees and shrubs used for groups and alleys.

The bark of magnolias are gray or brown, smooth, crusty, or flaky. Shoots with large ring traces and narrow ring marks from the stipules. The kidneys are large, narrowly conical or fusiform, with 1 or 2 scales. The leaves are large, mostly elliptical or obovate, entire, pinnate venous; second-order veins that do not reach the edge of the leaf. Stipules cover the young leaf.

Flowers homogeneous, usually very large, fragrant, white, cream or purple, solitary, terminal; Perianth of a three-leafed bowl, 6–9–12 petals, arranged in a circle of 2, 3 or 4, joined on top of each other. Stamens and pistils are numerous and are collected in a spindle-shaped, elongated vessel. The fruit is a conical folded leaf, the seeds are wedge-shaped, triangular, black, small embryos immersed in the oily endosperm, fleshy red, or pink seeds, the leaves hang on thin seed fibers when opened. Magnolia flowers are pollinated by beetles because they bloom before bees and butterflies appear. They do not contain nectar, but pollinating insects are attracted by a delicate sweet smell.

Magnolia seeds multiply for in the fall new cooked fruits collection and from them seeds get need seeds called sarkotesta solid shell with coated, many botanists from planting before take throw away advice they give. The shell softens for seeds 3–4 days during water with a spill throw need, then big sieve a how much times washed. That's it in a way processing given seeds immediately or the next year in February - March planted.

Seeds wet and cool in the environment save need, for example, a refrigerator middle shelf, moistened sphagnum moss wrapped and into placed. Plastic bag Mold appears to prevent getting for them fungicides with in advanced treatment possible. Prepared seeds, rule as closed 2 cm to the ground (boxes, buckets) in depth before they spilled into the grooves planted. Soil light, peat-sandy, lime without traces to be need Next year, seedlings 15-20 cm in height from then them separately in containers harvested, universal fertilizer with feeding possible.

Hot season grown magnolias open in the air, in the winter sea in a bright room to be stored possible, his temperature 5-10 ° C. And only of life third in the year branches constant to live for open to the ground transfer need Review magnolia as you stand multiplication seed method very much difficult From this except seeds obtained plants always of the parents type properties save will not remain.

Layer play Method

From the intersection new magnolia get much easy and this from seeds obtained samples than faster blooms. Layer - from parents inseparable of the plant root shot branch Reproduction for elasticity save the rest of the trap branches is selected. They are to the ground bent to the ground metal brackets with fastened and above soil with closes. Young sprouting place touch standing part of in the bark small a cut This is done method of the roots a little earlier appear to be opportunity gives Roots good developed then young tree from parents separated, permanent place planted.

Magnolia The plant is in Namangan conditions from the stem grow good the result does not give. No only cuttings multiplied plant unit but time passes with dry up remains. This plant from seed multiply gives good results. Because undamaged seeds slowly slowness with peat to the ground exacerbate it quickly unit comes out. Magnolia Uzbekistan a lot-grown type of this Magnolia grandiflora from seed good the result gives from the pen sea good the result get difficult. Magnolia The plant is in the climatic conditions of Namangan in greenhouses grown coming. Magnolia in greenhouses in the summer every day water pour stand up recommended are given. Because it is very watery demanding plant. Winter days while within 2 or 3 days times irrigation to the goal appropriate. A lot water injection as a result plant leaves quickly is dark enters and begins Summer Hot days up to 40 ° C heat resistant type is Cold days down to -3 ° C to the cold resistant plant.

Basic planted soil peat and biohumus. In some cases, to peat and biohumus less soil and fuzzy are added. Because the fuzzy composition is very rich in minerals. Magnolia The plant grows 20–30 cm in 1 year. Ours conditions more magnolia black disease does. It is a disease signs leaves darken the stay show will be. Such cases magnolia thirsty irrigation disease prevent known amount received will be.

REFERENCES

1. Misirova, S. A. "Systematic types of fungi of allocated and determined types from decorative flowers in conditions region Tashkent." *Agricultural sciences* 6.11 (2015): 1387.
2. Misirova, S. A. "Determining of the measure disease control ornamental crops during the growing season in the conditions Tashkent region." *Global Journal of Bio-Sciences and Biotechnology* 5.1 (2016): 119-124.
3. Abdumutalovna, Misirova Surayyo, and Sarimsaqova Nilufar Sobirjonovna. "Bioecology of Fungi-Pathogens of Flower Crops and the System to Combat Them." *Agricultural sciences* 7.08 (2016): 539.
4. MISIROVA, SA, and NN ERNAZAROVA. "FIGHTING MEASURES THE DISEASE CAUSES A VERY DANGEROUS FUNGAL SPECIES WIDESPREAD IN TASHKENT REGION." *International Journal of Botany and Research (IJBR)* 6 (2016): 5-12.
5. Misirova, Surayyo. "Technology of growing orchid flowers from seeds." *E3S Web of Conferences*. Vol. 390. EDP Sciences, 2023.
6. MISIROVA, SA. "TECHNOLOGY OF CULTIVATION AND REPRODUCTION OF ORNAMENTAL AND UNIQUE ORCHID FLOWER IN NAMANGAN CONDITIONS." *World Bulletin of Social Sciences* 17 (2022): 156-164.
7. Urmonovich, Numonov Otabek. "MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES." *ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ* 14.5 (2023): 3-5.
8. Misirova, S. A. "BIOLOGICAL CHARACTERISTICS OF FUNGAL SPECIES THAT CAUSE DISEASES OF ONION FLOWERS AND MEASURES TO COMBAT THEM." (2022).
9. Misirova, S., and M. Haydarova. "Flowers from Nederland are Considered to Develop in the Climatic Conditions of Uzbekistan and Are Identified the types of Fungus." *Annals of the Romanian Society for Cell Biology* (2021): 5922-5929.
10. Misirova, S. A., et al. "Determination types of fungi-pathogens of ornamental flower crops in conditions region Namangan." *ISJ Theoretical & Applied Science*, 10 (66) (2018): 185-189.
11. Misirova, S. A., M. U. Davlatova, and Sh O. Tuhtaboeva. "Biological Characteristics of Fungal Pathogens of Bulb Flowers and Control Measures." *JournalNX*: 207-214.
12. Misirova, S., et al. "Growing Dutch tulips in Namangan region." *Bulletin of Agrarian Science of Uzbekistan* 1 (2021).
13. Sabirov, Ravshan Z., et al. "Volume-sensitive anion channels mediate osmosensitive glutathione release from rat thymocytes." *PLoS One* 8.1 (2013): e55646.
14. Rashidovna, Melanova Nazira, and Numonov Otabek Urmonovich. "Comparative Characteristics of the Leaving of Glutathione From Cells of Different Types." *International Journal on Orange Technologies* 2.10: 79-82.
15. Melanova, N. R., M. U. Davlatova, and O. Numanov. "The Effect of Extracellular Glutathione on the Regulation of Thymocyte Volume in Rats under Conditions of

- Hypoosmotic Stress." Annals of the Romanian Society for Cell Biology (2021): 7032-7038.
16. Sabirov, R.Z., Kurbannazarova, R.S., Melanova, N.R. and Okada, Y., 2010, January. Swelling-induced release of glutathione from rat thymocytes. In JOURNAL OF PHYSIOLOGICAL SCIENCES (Vol. 60, pp. S13-S13). 1-11-11 KUDAN-KITA, CHIYODA-KU, TOKYO, 102-0073, JAPAN: SPRINGER TOKYO.
 17. Melanova, Nazira Rashidovna, and Sayyora Abduqahharovna Yulchiyeva. "EFFECT OF EXTRACELLUIAR GLUTATHIONE ON COLLOID-OSMOTIC LYSIS OF HUMAN RED BLOOD CELLS." Scientific and Technical Journal of Namangan Institute of Engineering and Technology 2.2 (2020): 144-149.
 18. Choriyeva, Nargiza Mamarajabovna, and Nazira Rashidovna Melanova. "STUDY OF LYSIS OF HUMAN ERYTHROCYTES UPON ADMINISTRATION OF GOSSYPOL, MEGOSIN AND BATRIDEN." Scientific and Technical Journal of Namangan Institute of Engineering and Technology 1.9 (2019): 55-58.
 19. Меланова, Назира Рашидовна. "Сравнительная характеристика выхода глутатиона из различных типов клеток." Universum: химия и биология 5 (59) (2019): 9-12.
 20. Насритдинов, Ахмаджон Абдухамидович, and Хусниддин Тургунбоевич Киргизов. "Агрегат для полосной обработки почвы." Современные научные исследования и инновации 12 (2015): 412-416.
 21. Байбобоев, Набижон Гуломович, et al. "Энергоресурсосберегающий комбинированный агрегат для обработки почвы." Вестник Рязанского государственного агротехнологического университета им. ПА Костычева 3 (23) (2014): 42-44.
 22. Насритдинов, А. А., and А. В. Рязанов. "Оптимальные условия установки углоснима." Техника в сельском хозяйстве 6 (2003): 34-35.
 23. Насритдинов, Ахмаджон Абдухамидович. "Результаты исследования формы лобовой поверхности стойки чизеля-культиватора." Universum: технические науки 1 (58) (2019): 18-20.
 24. Байбобоев, Набижон Гуломович, and Ахмаджон Насритдинов. "Теоретические определение перемещение частиц почвы по поверхности углоснима." Science Time 6 (18) (2015): 84-89.
 25. Misirova, S. A., M. U. Davlatova, and Sh O. Tuhtaboeva. "Biological Characteristics of Fungal Pathogens of Bulb Flowers and Control Measures." JournalNX: 207-214.
 26. Шамситдинов, Ф. "Результаты опыта." Защита и карантин растений 5 (2003): 27-27.
 27. Абдуалимов, Ш. Х., and Ф. Р. Шамситдинов. "Влияние применения стимуляторов роста на всхожесть семян, рост, развитие и урожайность хлопчатника в условиях светлых сероземных каменистых почв Наманганской области Республики Узбекистан." Актуальные проблемы современной науки 5 (2019): 47-51.

28. Абдуалимов, Шухрат Хамадуллаевич, and Фазлиддин Расулович Шамситдинов. "НАМАНГАН ВИЛОЯТИНИНГ ҚИР АДирЛИ ТОШЛОҚ ЕРЛАРИДА ЯНГИ СТИМУЛЯТОРЛАРНИНГ ҒЎЗА БАРГ ЮЗАСИ ВА ҲОСИЛДОРЛИГИГА ТАЪСИРИ." Журнал Биологии и Экологии 1 (2019).
29. Urmonovich N. O. MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES //ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ. – 2023. – Т. 14. – №. 5. – С. 3-5.
30. Rashidovna, Melanova Nazira, and Numonov Otabek Urmonovich. "Comparative Characteristics of the Leaving of Glutathione From Cells of Different Types." International Journal on Orange Technologies 2.10: 79-82.
31. Abdukhamidovich, Nasritdinov Akhmadzhon, and Numonov Otagek Urmonovich. "The Results of Theoretical Studies of the Chisel Cultivator Rack Frontal Surface Shape." Annals of the Romanian Society for Cell Biology (2021): 5930-5938.
32. Abdukhamidovich, Nasritdinov Akhmadzhon, Muhabbat Davlatova Urmanovna, and Numonov Otabek Urmonovich. "Strip Till Age of Soil for Deuteric Sowing (Second Crop)." International Journal on Orange Technologies 3.4 (2021): 71-74.
33. Abduhamidovich N. A. et al. MANGOSTIN DARAXTI VA MEVASINI TIBBIYOTDA FOYDALANISH //Journal of new century innovations. – 2023. – Т. 28. – №. 2. – С. 12-14.
34. Melanova, N. R., M. U. Davlatova, and O. Numanov. "The Effect of Extracellular Glutathione on the Regulation of Thymocyte Volume in Rats under Conditions of Hypoosmotic Stress." Annals of the Romanian Society for Cell Biology (2021): 7032-7038.
35. Рахманов, Дилшод Ортикбаевич, and Муҳаббат Ўрмоновна Давлатова. "ОРГАНОЛЕПТИЧЕСКИХ ОЦЕНКА СВОЙСТВ РЫБ И РЫБНЫХ ПРОДУКТОВ." O'ZBEKISTONDA FANLARARO INNOVATSIYALAR VA ILMIY TADQIQOTLAR JURNALI 2.14 (2022): 583-585.
36. Юсупова, Махпуза Нумановна. "БИОЛОГИЧЕСКИЙ МЕТОД ЗАЩИТЫ РАСТЕНИЙ." Scientific Impulse 1.9 (2023): 1460-1464.
37. Юсупова, Махпуза Нумановна. "АНОРНИ ЗАРАКУНАНДАЛАРДАН ҲИМОЯЛАШ." PEDAGOG 6.4 (2023): 562-567.
38. Юсупова, Махпуза Нумановна. "САБЗАВОТ ЭКИНЛАРИГА БИОЛОГИК КУРАШ ҲАҚИДА МУЛОХАЗАЛАР." Scientific Impulse 1.9 (2023): 1469-1473.
39. Юсупова, Махпуза Нумановна. "ФАРҒОНА ВОДИЙСИ ШАРОИТИДА ИГНА БАРГЛИ ДАРАХТЛАРНИ ЗАРАКУНАНДАЛАРДАН ҲИМОЯЛАШ." SO 'NGI ILMIY TADQIQOTLAR NAZARIYASI 6.4 (2023): 316-320.
40. O'rmonovna, Davlatova Muhabbat. "MANGOSTIN DARAXTI VA UNING KIMYOVIY XUSUSIYATLARI." INNOVATION IN THE MODERN EDUCATION SYSTEM 3.25 (2022): 1-4.
41. Юсупова, Махпуза Нумановна. "УФТ: 635 САБЗАВОТ ЭКИНЛАРИГА БИОЛОГИК КУРАШ ҲАҚИДА МУЛОХАЗАЛАР." Научный импульс: 355.

42. Qurbonov, I. "Tulip varieties imported from the netherlands technology of cultivation of namangan region. galaxy international interdisciplinary research journal (giirj) issn (E): 2347-6915 Vol. 9." (2021).
43. Kurbanov, I. G. "CARE OF TULIP VARIETIES OF THE NETHERLANDS IN THE CLIMATIC CONDITIONS OF THE NAMANGAN REGION." American Journal of Interdisciplinary Research and Development 6 (2022): 117-120.
44. Qurbonov I. E-RECRUITMENT: SOCIAL MEDIA AND RECRUITING //InterConf. – 2021.
45. Qurbonov, Ibragim Sharifjonovich. "CLONELY MICRO-CULTIVATION OF PLANTS AND ITS APPLICATION TO AGRICULTURE." Scientific and Technical Journal of Namangan Institute of Engineering and Technology 1.4 (2019): 74-78.
46. Юсупова М. Н., Ахмедова М. М. МЕВАЛИ ДАРАХТЛАРНИ ЗАРАРКУНАНДАЛАРИГА УЙЎНЛАШГАН КУРАШ ЧОРАЛАРИ //ЖУРНАЛ АГРО ПРОЦЕССИНГ. – 2020. – Т. 2. – №. 8.
47. Ходжаев, Ш. Т., Сагдуллаев, А. У., Исаев, О. Б., & Юсупова, М. Н. (2011). Проблемы защиты растений в Узбекистане. Защита и карантин растений, (8), 23-24. Юсупова М. Особенности защиты хлопчатника посеянного под пленки от вредных организмов //Автореф. канд. дисс./М. Юсупова–Ташкент. – 2001.
48. Ходжаев, Ш. Т., Юсупова, М. Н., Курызов, Ш., & Саттаров, Н. (2008). Перспективы биологической защиты хлопчатника от хлопковой совки. Сб. трудов.-Ташкент: Таллин, 44-49.
49. Yusupova M. N., Nosirov B. Z. Pests of cotton and straw control at collection //EPRA International Journal of Multidisciplinary Research (IJMR)-Peer Reviewed Journal. – 2020. – Т. 6. – №. 12. – С. 57-61.
50. Yusupova M. N., Axmedova M. M. Mevali daraxtlarni zararkunandalariga uygunlashgan kurash choralari //Jurnal JURNAL AGRO PROTSESSING. Data publikatsii. – 2020. – №. 8. – С. 12.
51. Yusupova M. N. Biological method of crop protection in the fergana valley //Agrarian science. – 2018. – №. 6. – С. 68-70.
52. Urmonovich, Numonov Otabek. "MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES." ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ 14.5 (2023): 3-5.
53. MN, Yusupova, and B. Z. Nosirov. "Control Of Cotton Pests On Stubble Lands." International Journal of Applied 10.2 (2015): 99-108.
54. Юсупова М. Н., Тургунова А. Н., Очилов С. Н. Система интегрированной защиты растений //Российский электронный научный журнал. – 2015. – №. 1. – С. 169-174.
55. Alimzhanova Z. I., Kadyrova D. S., Yusupova M. N. Ceramic pigments based on raw materials from Uzbekistan //Glass and Ceramics. – 2014. – Т. 70. – №. 11-12. – С. 441-443.

56. Yusupova M. N., Gapparov A. M. Biological Method Of Plant Protection In Uzbekistan //The American Journal of Agriculture and Biomedical Engineering. – 2020. – Т. 2. – №. 11. – С. 29-32.
57. Rashidovna M. N., Urmonovich N. O. Comparative Characteristics of the Leaving of Glutathione From Cells of Different Types //International Journal on Orange Technologies. – Т. 2. – №. 10. – С. 79-82.
58. Юсупова М. Н., Носиров Б. З. БИОЛОГИЧЕСКИЙ МЕТОД ЗАЩИТЫ РАСТЕНИЙ В УЗБЕКИСТАНЕ //Научно-практические пути повышения экологической устойчивости и социально-экономическое обеспечение сельскохозяйственного производства. – 2017. – С. 498-501.
59. Urmonovich, N. O. (2023). MANGOSTEEN NUTRITIONAL PRICE AND FUNCTIONAL PROPERTIES. ОБРАЗОВАНИЕ НАУКА И ИННОВАЦИОННЫЕ ИДЕИ В МИРЕ, 14(5), 3-5.
60. Yusupova M. et al. Protection of after harvest cultures-as a reservetors of cotton pests //Agriculture and Biology Journal of North America. – 2013. – Т. 4. – №. 5. – С. 576-582.
61. Ходжаев, Ш. Т., Юсупова, М. Н., Юлдашев, Ф., Исаев, О. Б., & Шокирова, Г. (2011). Борьба с вредителями хлопчатника на пожнивных культурах в севообороте. Вестник защиты растений, (2), 46-52.
62. Yusupova M. N. et al. Possibilities of the biological method of cotton plant protection //Agriculture and Biology Journal of North America. – 2011. – Т. 2. – №. 5. – С. 742-744.
63. Ходжаев, Ш. Т., Юсупова, М. Н., Юлдашев, Ф., & Жамалов, А. Г. (2010). Хлопковая совка на пожнивных культурах. Защита и карантин растений, (12), 22-23.
64. Хайдарова, Х. А., Юсупова, М. Н., Ихтиярова, Г. А., & Хайдаров, А. А. ПОЛУЧЕНИЕ ХИТОЗАНА ИЗ ПОДМОРА ПЧЕЛ APIS MILLIFERA. Сучасний рух науки: тези доп. XI міжнародної науково-практичної інтернет-конференції, 8-9 жовтня 2020 р.–Дніпро, 2020.–Т. 2.–426 с., 352.
65. Yusupova M., Turgunova A., Ochilov S. INTERGRATED PLANT PROTECTION SYSTEMS.